



Animal and Plant
Health Inspection
Service

Veterinary
Services

June 2026

Influenza A Virus in Swine Surveillance

Fiscal Year 2026 Quarterly Report

Surveillance Summary for First Quarter Fiscal Year 2026: October 1 to December 31, 2025

Report Summary

- This report covers the first quarter (Q1) of fiscal year (FY) 2026, from October 1 to December 31, 2025.
 - 1,095 samples were tested from 868 submitted accessions in Q1 FY 2026.
 - H1N2 was the predominant subtype reported in USDA data for the quarter.
 - Over the last 8 quarters, H1N1 was the predominant subtype in Regions 1, 3, and 5 and H1N2 was the predominant subtype in Regions 2 and 4.
 - The Agricultural Research Service (ARS) characterized 175 isolates with published sequences in GenBank by phylogenetic analysis.
 - The National Veterinary Services Laboratory's (NVSL) Diagnostic Virology Laboratory (DVL) provided 29 isolates to one academic and two government entities. NVSL received 30 isolates into the repository.

Key Points

- All IAV-S submissions are voluntary and based on clinical case submissions to veterinary diagnostic labs. These data are not a statistically representative sampling of the U.S. swine population.
- Due to the voluntary nature of this surveillance, the information in this report cannot be used to determine regional and/or national incidence, prevalence, or other epidemiological measures, but it may help identify IAV-S trends.
- The report provides data from both national and regional levels.
- Limited accessions from a region can skew data and lead to misinterpretation. Therefore, less inference can be applied to results from Regions 3, 4, and especially 5.
- Where relevant, this report includes previous years' data for historical perspective.

Introduction

This report, based on data received into the database as of April 7, 2026, provides a brief update on the status of national surveillance for influenza A virus in swine (IAV-S) for producers, swine practitioners, diagnosticians and the public. Summaries in this report may differ from those provided in past reports due to the regular addition of data from participating laboratories. The IAV-S surveillance program is voluntary and, as a result, the accessions and samples submitted represent a subset of the swine population. The surveillance system is not representative of the total U.S. domestic swine population; therefore, the data cannot be used to determine IAV-S prevalence or other epidemiologic measures in the swine population; however, the data may help identify influenza trends in swine. Program guidance indicates samples should only be collected from animals displaying influenza-like illness. Reporting months are based on the month the sample was collected. When the submitter does not report relevant information, data are recorded as "unknown."

A laboratory accession generally represents a set of samples collected at a single premises on a single day and subsequently received at the laboratory. While a nasal swab or lung tissue sample represents a single animal within the herd, a single oral fluid sample may represent one to two pens of animals in a herd. A positive sample status is based on the screening real-time reverse transcriptase polymerase chain reaction (rRT-PCR) on one or more samples within the accession. The subtype result is based on rRT-PCR-based subtyping assays. Virus isolation (VI) and sequencing in the National Animal Health Laboratory Network (NAHLN) labs are only attempted on rRT-PCR positives meeting criteria, with sequences deposited into [GenBank](#), the public sequence database. On a monthly basis, USDA NVSL also performs whole genome sequencing (WGS) on a selected subset of virus isolates received into the repository through the surveillance program and deposits those sequences into [Genbank](#). On a quarterly basis, a phylogenetic analysis is performed by USDA's Agricultural Research Service (ARS) National Animal Disease Center (NADC) influenza researchers; phylogenetic analyses are based on all successful USDA surveillance sequencing results deposited into [GenBank](#).

IAV-S Program Information

Current data, quarterly reports, and program information are available on the USDA APHIS IAV-S program website: <https://www.aphis.usda.gov/aphis/ourfocus/animalhealth/animal-disease-information/swine-disease-information/influenza-a-virus>

Requesting IAV-S Repository Isolates

Information on how to request available IAV-S repository isolates is available at: <https://www.aphis.usda.gov/sites/default/files/orderingiav-srepositoryisolates.pdf>

The full algorithm, including specific diagnostic parameters, timelines, and instructions for unusual results, is distributed to participating NAHLN laboratories through NVSL's Quality Management System.

Phylogenetic and genetic diversity data from the IAV-S surveillance system are visualized in the octoFLUshow interactive platform: <https://flu-crew.org>

IAV-S Surveillance Objectives

[USDA's National Surveillance Plan for Swine Influenza Virus in Pigs \(July 2010\)](#) describes the current surveillance system for IAV in swine in detail. The surveillance objectives are to:

1. Monitor genetic evolution of endemic IAV in swine to better understand endemic and emerging influenza virus ecology;
2. Make influenza isolates from swine available for research and establish a data management system to facilitate genetic analysis of these isolates and related information; and
3. Select proper isolates for the development of relevant diagnostic reagents, updated diagnostic assays, and vaccine seed stock products.

Objective 1. Monitoring Genetic Evolution of Endemic IAV in Swine to Better Understand Endemic and Emerging Influenza Virus Ecology

Objective 1 is met through the submission of diagnostic laboratory samples to the surveillance system, collection of the viruses that are isolated from the samples, and analysis of the hemagglutinin (HA) and neuraminidase (NA) sequences that are generated at the NAHLN laboratories. Each month, selected viruses undergo whole genome sequencing by the NVSL. Phylogenetic analysis of the genetic sequences submitted through the surveillance program is provided through an interagency agreement with the USDA's Agricultural Research Service (ARS) National Animal Disease Center (NADC).

National Surveillance Data Summary

The Cycle threshold (Ct) maximum value differs depending on the sample type. If lung/nasal samples have a Ct value of 25 or less and oral fluid samples have a Ct value of 20 or less, virus isolation and sequencing will be attempted. If there is something unique related to the virus, like it is causing high mortality, but the samples have higher than the established maximum Ct values, they will still enter the surveillance stream.

In Q1 FY2026, 1095 samples were tested from 868 submitted accessions (Figure 1). Figure 2 shows the overall trends in number of accessions submitted, rRT-PCR and VI positive accessions and subtyped accessions with a notable increase in accessions submitted and positive accessions beginning in August 2022 and continuing through Q2 FY2025 with a large spike from January 2024 through May 2024. The positivity rate, calculated by the number of positive accessions divided by the total number of accessions submitted for each month, shows seasonal fluctuations but has otherwise remained relatively consistent since 2022.




Figure 1. Number of IAV-S laboratory accessions and samples tested in swine FY2022 through Q1 FY2026

Figure 3 shows the number and distribution of subtype detections in Q1 FY2026. A total of 146 samples were subtyped, including H1N1 (n=42), H1N2 (n=59), H3N2 (n=40), and mixed subtype (n=5).

Figure 4 breaks down accessions by rRT-PCR subtype for FY2022 through Q1 FY2026. H1N1 was the predominant subtype detected in FY2022-FY2024 and H1N2 being the predominant subtype for FY2025 and Q1 FY2026. It is important to note that there is wide genetic diversity within each subtype.

Figure 5 displays the number of VI attempts in purple, successful VI attempts in orange, and the number of sequenced viral isolates submitted to [GenBank](#) in blue.

Influenza A Virus in Swine Surveillance Quarterly Report for Fiscal Year 2026, Quarter 1




Figure 2. IAV-S number of accessions submitted, subtyped accessions, rRT-PCR positive accessions, virus isolation positive accessions, and positivity rate over time (monthly), FY2022 through Q1 FY2026




Figure 3. Number of IAV-S subtype detections in Q1 FY2026




Figure 4. Breakdown of IAV-S accessions by subtype rRT-PCR from FY2022 through Q1 FY2026

Influenza A Virus in Swine Surveillance Quarterly Report for Fiscal Year 2026, Quarter 1




Figure 5. Number of IAV-S virus isolations attempted, positive virus isolations, and GenBank submissions from FY2022 through Q1 FY2026

Lab accessions were evaluated by age-class for the first quarter in FY2026. The most common subtype isolated in all of the classes was H1N2 (please note there were zero positive accessions from the sow/boar class) (Table 1). Table 2 displays the number of IAV-S positive accessions by specimen and viral subtype. Lung tissue yielded the highest percent of subtyped accessions in successful virus isolation (VI) attempts, with nasal swabs/wipes and oral fluids at 13% and 15% respectively. All sample types yielded at least 69% or above in percent of subtyped accessions sequenced and sent to GenBank, peaking at 88% for both lung and nasal swabs/wipes and oral fluids at 69%. Please note the low number of accessions with subtypes reported for nasal swabs/wipes and oral fluids.

Table 1. Number of positive accessions tested for IAV-S by age class and viral subtype, Q1 FY2026

Age Class (group)	Number of H1N1	Number of H1N2	Number of H3N1	Number of H3N2	Number of Mixed
Suckling	2	8	0	9	1
Nursery	14	17	0	16	3
Grow/Finish	14	17	0	4	1
Sow/Boar	0	0	0	0	0
Not Recorded/Unknown	4	4	0	3	0

Table 2. Number of positive accessions* tested for IAV-S by specimen type and by viral subtype, Q1 FY2026

Specimen Type (group)	Number of accessions with subtype reported	Percent of subtyped accessions with positive virus isolation	Percent of subtyped accessions sequenced and sent to GenBank	Number of H1N1	Number of H1N2	Number of H3N1	Number of H3N2	Number of Mixed	Number of samples sequenced and sent to GenBank
Lung	96	63%	88%	27	36	0	21	0	84
Nasal Swab or Wipe	8	13%	88%	0	4	0	3	0	7
Oral Fluids	13	15%	69%	3	5	0	1	0	9
Other Specimens	0	-	-	0	0	0	0	0	0

*Accessions may include samples with multiple specimen types. In these cases, individual accessions are counted in more than one specimen type category.

Regional surveillance data

In this section, we present IAV-S surveillance data across five different regions of the United States (Figure 6). These regions are based on former USDA administrative districts only and do not represent specific industry distributions. Submissions are voluntary, as is providing any identifying information beyond State of animal origin with the submission. Therefore, regional and/or national incidence, prevalence, or other epidemiological measures cannot be determined from this data.




Figure 6. A map of the regions for national IAV-S surveillance

Summary of Regional Data from ARS

Table 3 lists the predominant HA/NA phylo-type pairs by region from January 2025 through December 2025, with predominant being defined as comprising at least 10% of a region's HA/NA pairs. The total number column displays the total number of isolates that were phylo-typed for each region during that time. Historically, region 5 submits substantially fewer accessions than the other regions. Region 2 contains the most diversity of all the regions.

Figure 7 shows the distribution of rRT-PCR subtyped accessions across the five regions for Q1 FY2024 through Q1 FY2026. Over the last 8 quarters, H1N1 was the predominant subtype in Regions 1,3, and 5 (please note the overall low accessions for Region 5 (n=14) and H1N2 was the predominant subtype in Regions 2 and 4.

Table 3. Summary of predominant IAV-S HA/NA* phylo-types by US region for the 1-year window from January 2025 through December 2025

Region	Total Number	Predominant HA/NA subtypes
1	78	H1N1 (H1-1A.3.3.3-c3 / N1-C.3.2) (n=135) H3N2 (H3-2010.1 / N2-2002B) (n=133) H1N2 (H1-1B.2.1 / N2-1998B) (n=131)
2 ⁺	546	H1N1 (H1-1A.3.3.3-c3 / N1-C.3.2) (n=123) H1N2 (H1-1B.2.1 / N2-1998B) (n=96) H3N2 (H3-2010.1 / N2-2002B) (n=87)
3	69	H1N2 (H1-1A.1.1.3 / N2-2002A) (n=10) H1N2 (H1-1B.2.1 / N2-2002B) (n=9) H1N1 (H1-1A.3.3.3-c3 / N1-C.3.2) (n=8)
4	45	H1N2 (H1-1B.2.1 / N2-1998B) (n=15) H3N2 (H3-2010.1 / N2-2002B) (n=10) H1N1 (H1-1A.3.3.3-c3 / N1-P) (n=3)
5 ⁺⁺	4	H1N2 (H1-1A.1.1.3 / N2-2002B) (n=1) H1N1 (H1-1A.3.3.3-c3 / N1-P) (n=1) H3N2 (H3-2010.1 / N2-2002B) (n=1)
All	742	H3N2 (H3-2010.1 / N2-2002B) (n=148) H1N1 (H1-1A.3.3.3-c3 / N1-C.3.2) (n=146) H1N2 (H1-1B.2.1 / N2-1998B) (n=138)

*HA/NA pairs included if they comprise over 10% from a region

⁺ Most diversity of all regions

⁺⁺ Low participation




Figure 7. Distribution of IAV-S rRT-PCR subtyped accessions across five US regions for Q1 FY2024 through Q1 FY2026

Phylogenetic analysis of sequences from the IAV-S surveillance system

Phylogenetic analysis of gene sequences of IAV in swine is conducted by ARS to further examine the genetic changes that occur in HA and NA genes of this rapidly changing virus. Through collaboration with ARS, a dataset of 175 isolates with published sequences in [GenBank](#) was characterized by phylogenetic analysis for the Q1 FY2026 report. This analysis provides information on the genetic diversity and evolutionary patterns of IAV in swine and allows for inferences about population and/or vaccine immunity.¹

The following series of bar charts parse the data into an approximately 2-year window by quarters and region, describing virus subtypes (Figure 8) and phylogenetic clades of H1, H3, N1 and N2 subtypes (Figures 9-12). Regional charts depicting various combinations of HA and NA are available in Appendix 1.

Figure 8 demonstrates the four subtypes H1N1, H1N2, H3N1 and H3N2 across the five regions. Region 2 reported the most submissions with Regions 1, 3, and 4 all reporting a similar number of submissions. All were a mixture of mostly H1N1, H1N2, and H3N2 subtypes. Limited accessions from a region can skew data and lead to misinterpretation, therefore less inference can be applied to results from region 5.




Figure 8. Temporal distribution of Influenza A virus of swine subtype by region for Q2 FY2024 to Q1 FY2026

¹ The ARS dataset is comprised of IAV-S surveillance isolate sequences from Genbank. This represents only a subset of the complete IAV-S surveillance dataset that includes PCR diagnostic test-based results as well as sequencing results. Therefore, ARS dataset results, such as subtype percentages, differ from the complete IAV-S dataset results provided in other sections of this report.

National phylogenetic HA gene information

Due to the US clade classification being too coarse and no longer informative, starting in Q1 of FY2024, the HA genes from H1 subtype viruses are no longer classified using the US phylogenetic clades alpha, beta, gamma, delta1, delta2, or pandemic H1N1 2009 (H1N1pdm09). Similarly, for H3 subtype viruses, US phylogenetic clades Cluster IV, Cluster IV-A, Cluster IV-B, Cluster IV-C, Cluster IV-D, Cluster IV-E, Cluster IV-F, 2010.1, 2010.2, or human-like are no longer used. Instead, they are classified using global clades as described in these two published nomenclature systems: [A Phylogeny-Based Global Nomenclature System \(2016\)](#) and [Swine Influenza A Viruses and the Tangled Relationship with Humans \(2021\)](#). In Q1 FY2026, there were 6 H1 and 5 H3 clades detected. The predominant H1 HA clades were H1-1A.1.1.3, H1-1B.2.1, and H1-1A.3.3.3-c3, representing 78.7% of the H1 collection (n=150) (Figure 9). The predominant H3 HA clade was H3-2010.1, representing 68.0% of the H3 collection (n=25) (Figure 10). There was an increase of: 2.5% in clade H1-1A.3.3.3-c3, 1.8% in clade H1-1B.2.2, and 1.0% in clade H1-1A.3.3.2 over the last year. There was a decrease of: 3.7% in clade H3-1990.4.a, 1.3% in H3-1990.4.b1, and 0.8% in clade H3-2010.2 over the last year.




Figure 9. Temporal distribution of IAV-S H1 phylogenetic clades by region for Q2 FY2024 to Q1 FY2026




Figure 10. Temporal distribution of IAV-S H3 phylogenetic clades by region Q2 FY2024 to Q1 FY2026

National phylogenetic NA gene information

In Q1 FY2026, N1-C.3.2 and N1-P, were the predominant N1 phylogenetic-clades, representing approximately 95.0% of the N1 collection (n=60). In Q1 FY2026, the predominant N2 phylogenetic-clades were N2-2002B and N2-1998B, representing approximately 89.6% of the N2 collection (n=115). Figures 11 and 12 show the temporal distribution of IAV-S N1 and N2 phylogenetic-clades by region for Q2 FY2024 through Q1 FY2026.




Figure 11. Temporal distribution of IAV-S N1 phylogenetic-clades by region for Q2 FY2024 to Q1 FY2026




Figure 12. Temporal distribution of IAV-S N2 phylogenetic-clades by region for Q2 FY2024 to Q1 FY2026

Representative HA genes

Six months of IAV-S data, July 2025–December 2025, were used by the NADC to identify circulating HA clades. An objective algorithm was used to identify the best-matched field strain housed in the USDA IAV-S virus repository (Table 4).

Table 4. IAV-S Surveillance NADC Representative HA genes*

Strain	GenBank	Global Clade
A/swine/Missouri/A02858730/2025	PX114458	H1-1A.1.1.3
A/swine/Minnesota/A02980710/2025	PX596405	H1-1A.3.3.2
A/swine/Iowa/A02858735/2025	PX114466	H1-1A.3.3.3-c3
A/swine/Iowa/A02859054/2025	PZ019711	H1-1B.2.1
A/swine/Iowa/A02858843/2025	PX459590	H1-1B.2.2.2
A/swine/Minnesota/A02980758/2025	PX934434	H3-1990.4.a
A/swine/North_Carolina/A02858783/2025	PX368848	H3-2010.1
A/swine/Iowa/A02858897/2025	PX504850	H3-2010.1

*6-month HA1 objective algorithm and best-matched field strain in the repository was identified.

- July 2025 to December 2025 USDA HA data downloaded ($n = 264$ H1, $n = 68$ H3) and a phylogenetic tree was inferred. For each HA clade, an objective representative selection was made using PARNAS (<https://github.com/flu-crew/parnas>).
- The 5 H1 selections cover 85% of observed diversity; 3 H3 selections cover 54% of observed diversity.
- Clades were required to have a detection rate of at least 2% to be considered for selection ($n \geq 7$).
 - Omitted H1-1A.3.3.3-c1 ($n=3$), H3-1990.4.b1 ($n=1$), H3-1990.4.i ($n=5$), H3-2010.2 ($n=3$), and H3-2020.2 ($n=2$)
- Note that 2 selections were made within H3-2010.1, which represents 86% of all H3 detections.

Representative NA genes

Six months of IAV-S data, July 2025–December 2025, were used by the NADC to identify circulating NA clades. An objective algorithm was used to identify the best-matched field strain housed in the USDA IAV-S virus repository (Table 5).

Table 5. IAV-S Surveillance NADC Representative NA genes*

Strain	GenBank	Global Clade
A/swine/Iowa/A02859047/2025	PX985999	N1-C.3.2
A/swine/Indiana/A02858874/2025	PX480116	N1-P
A/swine/Iowa/A02858879/2025	PX480124	N1-C.2.1
A/swine/Iowa/A02858726/2025	PX114451	N2-1998B
A/swine/Montana/A02858889/2025	PX504837	N2-2002A
A/swine/Iowa/A02858770/2025	PX310588	N2-2002B

*6-month NA1 objective algorithm and best-matched field strain in the repository was identified.

- July 2025 to December 2025 USDA NA data downloaded ($n = 115$ N1, $n = 217$ N2) and a phylogenetic tree was inferred. For each HA clade, an objective representative selection was made using PARNAS (<https://github.com/flu-crew/parnas>).
- The 3 N1 selections cover 82% of observed diversity; the 3 N2 selections cover 61% of observed diversity.
- Clades were required to have a detection rate of at least 2% to be considered for selection ($n \geq 7$).
 - Omitted N2-2002 ($n=5$), N2-2016 ($n=3$), N2-LAIV-98 ($n=2$).

Gene constellations

Please note that complete WGS data and associated figures for Q1 FY2026 are not included in this report. This is due to a recent change in data processing timelines which impacted the availability of the most recent data needed for our comprehensive analysis. The comprehensive WGS analysis covering this time period will be provided in subsequent reports.

Objective 2. Make Influenza Isolates from Swine Available for Research and Establish a Data Management System to Facilitate Genetic Analysis of these Isolates and Related Information

A primary goal of the IAV swine surveillance program is to share selected virus isolates obtained through the surveillance system with public health, animal health, and academic researchers to facilitate genetic analysis and research on viruses of interest. The NVSL Diagnostic Virology Laboratory maintains a repository of the viruses submitted into the surveillance system and provides these viruses upon request.

In Q1 FY2026, the NVSL Diagnostic Virology Laboratory provided 29 isolates to one academic and two government entities. NVSL received 30 isolates into the repository. Table 6 reports the total number of virus isolates received into the repository each year from FY2014 through Q1 FY2026. Table 7 reports the total number of isolates by subtype available in the repository for sharing.

IAV-S isolates can be requested from the NVSL repository by following the instructions found at: <https://www.aphis.usda.gov/sites/default/files/orderingiav-srepositoryisolates.pdf>

Table 6. IAV-S isolates received in NVSL repository by fiscal year

Fiscal Year	Number of Isolates
Q1 FY2026	30
FY2025	1,091
FY2024	1,083
FY2023	1,035
FY2022	641
FY2021	1,108
FY2020	1,074
FY2019	1,055
FY2018	994
FY2017	844
FY2016	1,046
FY2015	883
FY2014	765
TOTAL	11,649

Table 7. Total number of subtyped IAV-S isolates collected from 2009- Q1 FY2026 and available through the NVSL repository

Subtype	Number of Isolates Received in Q1 FY2026	Total Number of Isolates in Repository
H3N2	5	3,745
H3N1	0	33
H1N1	11	4,350
H1N2	14	4,249
Mixed	0	304
TOTAL	30	12,681

Objective 3. Select Proper Isolates for Development of Relevant Diagnostic Reagents, Updating Diagnostic Assays, and Vaccine Seed Stock Products

USDA makes IAV-S isolates available in the public domain for further research. ARS-NADC conducts research on isolates obtained from the repository and sequences generated from the surveillance system. Genetic sequencing from the USDA program that is reported to [GenBank](https://www.ncbi.nlm.nih.gov/genbank/) is available for private corporations, government entities, academia, and other scientific community partners for research and vaccine strain selection and efficacy testing. NVSL and ARS staff are consulted as subject matter experts when necessary.

Conclusion


The IAV-S voluntary surveillance system in swine continues to provide insight into the genetic makeup of circulating influenza A virus in limited populations of commercial pigs. Genetic information and virus isolates are made publicly available for further research and possible vaccine strain selection and efficacy testing. Influenza A virus in swine remains a dynamic virus with high levels of genetic variability in the hemagglutinin and neuraminidase genes.

Appendix 1. Regional Charts of IAV-S HA and NA Combinations by Percentage

The following charts present the percentages of combinations of IAV-S HA and NA by region based on ARS-NADC phylogenetic analyses. The results are reported from January 2025 through December 2025 for regions 1-3, January 2025 through November 2025 for region 4, and January 2025 through October 2025 for region 5. These “heat maps” represent the percentage of combinations by using a color gradient where a deeper gradient color represents a greater percentage occurrence for a particular HA-NA combination. HA clusters are listed on the left vertical axis of the chart and NA clusters are listed on the bottom horizontal axis. Line up the HA cluster with the corresponding NA cluster to determine the occurrence of that particular combination.



Region 1



Total HA & NA combinations – 78



Region 2

Percentage of HA and NA combinations - Jan 2025 to Dec 2025


HA Clade	N1.C.2.1	N1.C.3.1	N1.C.3.2	N1.P	N2.1998B	N2.2002	N2.2002A	N2.2002B	N2.2016	N2.LAIV-98	HA. Total
H1.1A.1.1.3	2		1.6	4.2	8.2		0.5	2.6			19.1
H1.1A.3.3.2				5.7				0.2			5.9
H1.1A.3.3.3-c1	1.1							0.5			1.6
H1.1A.3.3.3-c3		0.5	22.5								23
H1.1B.2.1					17.6			2			19.6
H1.1B.2.2.2					2.2		1.3	4		0.2	7.7
H3.1990.4								0.2			0.2
H3.1990.4.a								2.6			2.6
H3.1990.4.i						1.3					1.3
H3.2010.1			0.2				0.9	15.9			17
H3.2010.2									1.3		1.3
H3.2020.2								0.5			0.5
NA.Total	3.1	0.5	24.3	9.9	28	1.3	2.7	28.5	1.3	0.2	
	N1.C.2.1	N1.C.3.1	N1.C.3.2	N1.P	N2.1998B	N2.2002	N2.2002A	N2.2002B	N2.2016	N2.LAIV-98	HA. Total

NA Clade

Total HA & NA combinations – 546



Region 3



Total HA & NA combinations – 69



Region 4


Percentage of HA and NA combinations - Jan 2025 to Nov 2025

HA Clade	N1.C.3.2	N1.P	N2.1998B	N2.2002	N2.2002A	N2.2002B	HA.Total
H1.1A.1.1.3		2.2	2.2		4.4	6.7	15.5
H1.1A.3.3.2		6.7				2.2	8.9
H1.1A.3.3.3-c3	2.2						2.2
H1.1B.2.1			33.3			4.4	37.7
H3.1990.4				2.2			2.2
H3.1990.4.b1						4.4	4.4
H3.1990.4.i				4.4			4.4
H3.2010.1			2.2			22.2	24.4
NA.Total	2.2	8.9	37.7	6.6	4.4	39.9	
	N1.C.3.2	N1.P	N2.1998B	N2.2002	N2.2002A	N2.2002B	HA.Total
	NA Clade						

Total HA & NA combinations – 45



Region 5



Total HA & NA combinations – 4